

This article was downloaded by: [Universitetsbiblioteket i Bergen]

On: 22 April 2015, At: 00:29

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Journal of the American Statistical Association

Publication details, including instructions for authors and subscription information:
<http://www.tandfonline.com/loi/uasa20>

On the Relative Importance of Input Factors in Mathematical Models

Andrea Saltelli^a & Stefano Tarantola^a

^a Andrea Saltelli and Stefano Tarantola are researchers at the Joint Research Centre of the European Commission, Institute for the Protection and Security of the Citizen, Ispra, Italy.

Published online: 31 Dec 2011.

To cite this article: Andrea Saltelli & Stefano Tarantola (2002) On the Relative Importance of Input Factors in Mathematical Models, *Journal of the American Statistical Association*, 97:459, 702-709, DOI: [10.1198/016214502388618447](https://doi.org/10.1198/016214502388618447)

To link to this article: <http://dx.doi.org/10.1198/016214502388618447>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

On the Relative Importance of Input Factors in Mathematical Models: Safety Assessment for Nuclear Waste Disposal

Andrea SALTELLI and Stefano TARANTOLA

This article deals with global quantitative sensitivity analysis of the Level E model, a computer code used in safety assessment for nuclear waste disposal. The Level E code has been the subject of two international benchmarks of risk assessment codes and Monte Carlo methods and is well known in the literature. We discuss the Level E model with reference to two different settings. In the first setting, the objective is to find the input factor that drives most of the output variance. In the second setting, we strive to achieve a preestablished reduction in the variance of the model output by fixing the smallest number of factors. The emphasis of this work is on how to define the concept of importance in an unambiguous way and how to assess it in the simultaneous occurrence of correlated input factors and non-additive models.

KEY WORDS: Analysis of variance; Correlated input; Nonadditive model; Sensitivity analysis.

1. INTRODUCTION

The work focuses on data that are the output from a computer code. Statisticians have concerned themselves with these kinds of datasets since the early 1990s (Sacks, Schiller, and Welch 1989; Sacks, Welch, Mitchell, and Wynn 1989b; Welch et al. 1992) addressing the issue from several viewpoints: (a) how to approximate the output from a complex computer code; (b) how to optimally design exploratory points to do (a); (c) how to characterize the empirical distribution of the output given probability distributions of the input; and (d) how to assess the importance of input factors in relation to (c). In this work, the emphasis is substantially on (d).

In recent years, global quantitative sensitivity analysis techniques have received considerable attention in the literature (RESS 1997; JSCS 1997; CPC 1999; JMCDA 1999). Many techniques have been developed that can be applied even to nonlinear, nonmonotonic models. In particular, in an earlier work (Saltelli, Tarantola, and Chan 1999) we proposed a model-free method for efficient global sensitivity analysis for sets of noncorrelated input factors. A review of applications of the same methods was given by Saltelli, Tarantola, and Campolongo (2000). Saltelli, Chan, and Scott (2000) presented a broad spectrum of techniques.

In the present work we extend the methodology for use in the simultaneous presence of correlated inputs and nonadditive models, and apply it to a well-known test case, the Level E model. The Level E test case was first used as a benchmark to compare the performance of computer codes used in safety assessment for nuclear waste disposal (OECD 1989), and later was used as a test model for sensitivity analysis techniques (OECD 1993). In this latter study, called Level S, the Level E model was run in a Monte Carlo fashion, with the model fed with samples from the distribution of the inputs. The Level S aimed to rank the uncertain input factors in order of importance according to two different criteria:

a. Reduction in the spread of the output at given times for a 5% reduction at each end of the input factor's range, that is,

taking a new distribution for that factor running between the former 5th and 95th percentiles

b. Change in the mean of the output at given times for a 5% increase of the central value of the factor's distribution, that is, shifting the whole distribution up, so that the bottom end is at the former 5th percentile.

The results showed that the sensitivity analysis methods applied in the Level S exercise were of limited value in ranking the importance of the input factors in the Level E test model. Depending on the formulation of the questions in Level S, the results were in fact dependent on the criterium selected (i.e., shrinking or shifting), and the effect of nonmonotonicity in the input-output relationship was not captured.

The main difficulty in sensitivity analysis at the time of the Level S exercise was on how to interpret its results. Although both criteria (a) and (b) are legitimate in principle, and answerable by straightforward Monte Carlo methods (OECD 1993), one does not know what question is being answered by a ranking of the input factors based on (a) or (b). Should we prioritize research on a given factor if it scores first in importance under (a)? Or under (b)? What if the answer depends on the percentage values previously stipulated? Should the same percentage apply to all factors, or should it include consideration of the likely reduction in uncertainty subsequently achievable in the laboratory, or in the field, for each individual factor? Was neglecting the covariance structure of the problem legitimate, or did it induce too large an error? A number of articles that followed the Level S report attempted to answer these questions (see Sec. 2.2).

In this work, we attempt to define "importance" in a rigorous fashion. The Level S benchmark revealed quite a latitude of different understandings of the concept of "importance" when applied to sensitivity analysis. We revisit the nonlinear, nonadditive Level E case study, complementing it with the input correlation that was omitted in the original exercise. As we show, a sensitivity analysis in the simultaneous occurrence

Andrea Saltelli (E-mail: andrea.saltelli@jrc.it) and Stefano Tarantola (E-mail: stefano.tarantola@jrc.it) are researchers at the Joint Research Centre of the European Commission, Institute for the Protection and Security of the Citizen, Ispra, Italy.

of correlation (a property of the input sample) and interaction (a property of nonadditive models) is nontrivial.

2. THE LEVEL E MODEL

2.1 Description of the Model

Level E was used both as a benchmark of Monte Carlo computation (Robinson and Hodgkinson 1987; OECD 1989) and as a benchmark for sensitivity analysis methods (Level S), (OECD 1993). The model predicts the radiologic dose to humans over geologic time scales due to the underground migration of radionuclides from a nuclear waste disposal site. The scenario considered in the model tracks the one-dimensional migration of four radionuclides (^{129}I and the chain $^{237}\text{Np} \rightarrow ^{233}\text{U} \rightarrow ^{229}\text{Th}$) through two geospheric layers with different hydrogeologic properties. The processes considered in the model are radioactive decay, dispersion, advection, and chemical reaction between the migrating nuclide and the porous medium. The repository is represented as a point source. Sometime after the steel canister containing the waste has lost its integrity (with the time of containment failure indicated by T), the release of radionuclides to the geosphere depends on only the leach rates ($k_{(i)}$) and the initial inventory ($C_{(i)}$). The source term for ^{129}I is given by

$$\begin{aligned} \frac{\partial C_I}{\partial t} &= -\lambda_I C_I, & t \leq T, \\ \frac{\partial C_I}{\partial t} &= -\lambda_I C_I - k_I C_I, & t > T, \end{aligned} \quad (1)$$

where C_I (mols) is the amount of ^{129}I and λ_I (yr^{-1}) and k_I (yr^{-1}) are the decay rate and the leaching rate for ^{129}I . The initial condition is $C_I(t=0) = C_I^0$, that is, the amount of ^{129}I at the time of vault closure (Table 1). The source term for ^{237}Np , the first element of the chain, is also described by (1),

in which the parameters λ_{Np} and k_C are used. (Here k_C is the leaching rate for the radionuclides of the chain.) The source term for ^{233}U is given by

$$\begin{aligned} \frac{\partial C_U}{\partial t} &= -\lambda_U C_U + \lambda_{Np} C_{Np}, & t \leq T, \\ \frac{\partial C_U}{\partial t} &= -\lambda_U C_U + \lambda_{Np} C_{Np} - k_C C_U, & t > T. \end{aligned} \quad (2)$$

The source term for ^{229}Th is described similarly by (2), where U is replaced by Th and Np is replaced by U . The migration through the geosphere is the core of the model. For example, the migration of ^{233}U is governed by

$$R_U^{(k)} \frac{\partial F_U^{(k)}}{\partial t} = v^{(k)} d^{(k)} \frac{\partial^2 F_U^{(k)}}{\partial x^2} - v^{(k)} \frac{\partial F_U^{(k)}}{\partial x} - \lambda_U R_U^{(k)} F_U^{(k)} + \lambda_{Np} R_{Np}^{(k)} F_{Np}^{(k)}, \quad (3)$$

where U stands for the isotope ^{233}U , Np stands for ^{237}Np , (k) refers to geosphere layer number k (1 or 2), R_i is the retardation coefficient for nuclide i (dimensionless), $F_i(x, t)$ is the flux (amount transported per unit time) of nuclide i in the geosphere at position x and time t (mols/yr), $v^{(k)}$ is the water travel velocity in the k th geosphere layer (m/yr), $d^{(k)}$ is the dispersion length in the k th geosphere layer (m), and λ_i is the decay constant of nuclide i (yr^{-1}).

The same equation holds for ^{229}Th provided that the index U is replaced by Th and the index Np by U . To simplify the model structure, the retardation coefficients R_U , R_{Np} , and R_{Th} were replaced in the Level E exercise by a single parameter R_C . The equation for ^{129}I is

$$R_I^{(k)} \frac{\partial F_I^{(k)}}{\partial t} = v^{(k)} d^{(k)} \frac{\partial^2 F_I^{(k)}}{\partial x^2} - v^{(k)} \frac{\partial F_I^{(k)}}{\partial x} - \lambda_I R_I^{(k)} F_I^{(k)}. \quad (4)$$

Table 1. List of Input Factors for the Level E Model

Notation	Definition	Distribution	Range	Units
T	Containment time	Uniform	[100, 1000]	yr
k_I	Leach rate for iodine	Log-uniform	$[10^{-3}, 10^{-2}]$	mols/yr
k_C	Leach rate for Np chain nuclides	Log-uniform	$[10^{-6}, 10^{-5}]$	mols/yr
$v^{(1)}$	Water velocity in geosphere's 1st layer	Log-uniform	$[10^{-3}, 10^{-1}]$	m/yr
$l^{(1)}$	Length of geosphere's 1st layer	Uniform	[100, 500]	m
$R_I^{(1)}$	Retention factor for I (1st layer)	Uniform	[1, 5]	—
$R_C^{(1)}$	Factor to compute retention coefficients for Np (1st layer)	Uniform	[3, 30]	—
$v^{(2)}$	Water velocity in geosphere's 2nd layer	Log-uniform	$[10^{-2}, 10^{-1}]$	m/yr
$l^{(2)}$	Length of geosphere's 2nd layer	Uniform	[50, 200]	m
$R_I^{(2)}$	Retention factor for I (2nd layer)	Uniform	[1, 5]	—
$R_C^{(2)}$	Factor to compute retention coefficients for Np (2nd layer)	Uniform	[3, 30]	—
W	Stream flow rate	Log-uniform	$[10^5, 10^7]$	m^3/yr
C_I^0	Initial inventory for ^{129}I	Constant	100	mols
C_{Np}^0	Initial inventory for ^{237}Np	Constant	1000	mols
C_U^0	Initial inventory for ^{233}U	Constant	100	mols
C_{Th}^0	Initial inventory for ^{229}Th	Constant	1000	mols
w	Water ingestion rate	Constant	.73	m^3/yr
β_I	Ingestion-dose factor for ^{129}I	Constant	56	Sv/mols
β_{Np}	Ingestion-dose factor for ^{237}Np	Constant	6.8×10^3	Sv/mols
β_U	Ingestion-dose factor for ^{233}U	Constant	5.9×10^3	Sv/mols
β_{Th}	Ingestion-dose factor for ^{229}Th	Constant	1.8×10^6	Sv/mols

The same equation holds for ^{237}Np provided that the index I is replaced by Np . Modeling of the biosphere is extremely simplified; via an ingestion factor and the water consumption rate, the dose to the most exposed individual of a hypothetical critical group is computed. The radiologic dose (measured in Sv/yr) from nuclide i is given by

$$\text{dose}_i(t) = \beta_i \frac{w}{W} F_i^{(2)}(I^{(2)}, t), \quad i = {}^{129}\text{I}, {}^{237}\text{Np}, {}^{233}\text{U}, {}^{229}\text{Th}, \quad (5)$$

where β_i is an ingestion-dose conversion factor and is assumed fixed, $F_i^{(2)}(I^{(2)}, t)$ is the flux at the end of the second layer (the output to the biosphere), w denotes the drinking water requirement for an individual in the most exposed critical group, and W is the stream flow rate. The quantity of interest in this study is the annual radiologic dose due to the four radionuclides,

$$Y(t) = \sum_i \text{dose}_i(t). \quad (6)$$

The overall predictive uncertainty about that dose is due to uncertainties in model parameters, both intrinsic (or stochastic), such as the time of canister failure, and due to our poor knowledge of the system, such as a poorly known kinetic parameter. The simulation model includes 12 uncertain input factors, listed in Table 1 together with a set of parameters that are assumed constant.

The probability distributions for each factor were selected on the basis of expert judgement. Such data refer to the original formulation of Level E (OECD 1989). Although we were aware of the existence of correlations (e.g., between R_I and R_C for each layer of the geosphere), these were omitted in the original system specifications for the sake of simplicity. For a subset of our simulations, we have reinstated these correlations by assuming that a serious vault failure induces high leaching for both iodine and the chain elements, and vice versa. Also, high release coefficients for iodine should be accompanied by high release coefficients for the chain elements, within a given geosphere layer. The geochemical properties in the two layers of the geosphere could hardly be independent from each other, and the water flows in the two layers should show a certain degree of dependency. Besides, the time of containment failure is likely correlated to the flow in the first layer, because corrosion will be faster if the flow is faster. The proposed correlation pattern is given in Table 2. The set of correlation values was defined by consulting with the authors of the benchmark (P. C. Robinson 2000, personal communication).

Table 2. Configuration for Correlated Input of the Level E Model

Pairs of correlated factors	Correlation
k_I, k_C	.5
$R_I^{(1)}, R_C^{(1)}$.3
$R_I^{(2)}, R_C^{(2)}$.3
$T, v^{(1)}$	-.7
$v^{(1)}, v^{(2)}$.5
$R_I^{(1)}, R_I^{(2)}$.5
$R_C^{(1)}, R_C^{(2)}$.5

To replicate the Level E exercise, the system (1)–(6) needs to be solved numerically. We used the Crank–Nicholson method (see Crank 1975; Prado, Homma, and Saltelli 1991). As an alternative, the system can be solved using an algorithm for accurate numerical inversion of the solution in the Laplace space to obtain the solution in the real space (Robinson and Hodgkinson 1987). We tackled below both cases (correlated and noncorrelated) using two alternative definitions of importance.

2.2 Previous Work on Level E

Following the Level S benchmark, a number of authors attempted to address the concept of importance in sensitivity analysis using the Level E as a key test case. Saltelli and Homma (1992) suggested an early version of the first-order conditional variance V_j (Hora and Iman 1989; Ishigami and Homma 1990) as a sensitivity analysis tool for nonmonotonic models. We treat this measure in Section 3.

Saltelli, Andres, and Homma (1993) improved the estimation procedure for V_j and applied it to the Level E example. The results are reported in Section 5.1. A version of V_j based on the ranks of the output values, rather than the values themselves, was also suggested to reduce the relative error of the estimates. It was realized later (Saltelli and Sobol' 1995), using, among others, the Level E case, that the improved reproducibility of the rank-based measure came at the cost of substantial alteration of the sensitivity pattern of the model output Y considered. The sensitivity pattern of "rank of Y " was substantially different from the sensitivity pattern of Y .

Level E was instrumental in the development of the total sensitivity indices S_{Tj} (Homma and Saltelli 1996), described in Section 4.1. In brief, the fact that some interaction terms were important for this model led to the development of a sensitivity measure incorporating both the first-order and higher-order effects of a given factor on model output.

The Level E proved instructive in other works. Saltelli and Sobol' (1995) used Sobol' quasi-random LP- τ sequences to sample the input factors space when computing conditional variances using the method of Sobol' (Sobol' 1990). Saltelli and Bolado (1998) compared the Sobol' and FAST methods to compute sensitivity measures. Saltelli et al. (1999) extended the FAST method to compute the total effect indices S_{Tj} . Draper et al. (1999) and Draper, Saltelli, Tarantola, and Prado (2000) used an extended version of the Level E to study the propagation of parametric and scenario uncertainty in a Bayesian framework. Saltelli, Tarantola, and Campolongo (2000) used different models, including Level E, to illustrate the effectiveness of variance based sensitivity measures.

3. LOTTERY SETTING 1

How do we judge the relative importance of model input factors? We assume to have information about the factors' probability distribution, either joint or marginal, with or without correlation, and that this knowledge comes from measurements, estimates, expert opinion, physical bounds, output from simulations, analogy with factors for similar species, and so forth.

We also assume as good practice in model use that from the many output variables generated by the model, we select

a single top-level variable, Y , that summarizes the thesis that the model is supposed to prove or disprove (Saltelli, Tarantola, and Campolongo 2000). Examples of Y include the extent of a geographic area where the concentration of a given pollutant exceeds a given threshold value, the estimated failure probability of a given system in a given time range, and the risk that the return of a given portfolio, falls below a preestablished value within a given time span. As a general rule, the functional of interest is selected so that it is meaningful with respect to the question asked of the model.

Let us assume that we can extract a sample from the probability distribution of the input factors and that we can compute the model output for every sample point. Also assume that each factor has a true, albeit unknown, value. We are hence poised to stipulate the following setting.

Lottery Setting 1. We are asked to bet on the factor that, if determined (i.e., fixed to its true value), would lead to the greatest reduction in the variance of Y .

The factors could be ranked according to $V(Y|X_i = x_i^*)$ or, equivalently, to $V(Y|X_i = x_i^*)/V$, where V is the unconditional variance of Y and $V(Y|X_i = x_i^*)$ is the variance obtained by fixing X_i to its true value x_i^* . The problem is that x_i^* is unknown for each X_i .

It would sound sensible to bet on the factor with the smallest value of $E[V(Y|X_i)]$, that is, the average of the conditional variance $V(Y|X_i = x_i^*)$ evaluated over all possible values x_i^* of X_i . Given that

$$V = V[E(Y|X_i)] + E[V(Y|X_i)], \quad (7)$$

betting on the lowest $E[V(Y|X_i)]$ is equivalent to betting on the highest $V[E(Y|X_i)]$.

Unsurprisingly, many practitioners of sensitivity analysis have come up with different estimates of $V[E(Y|X_i)]$ (in short, V_i) as a measure of sensitivity. Some have called this the *importance measure* (see Chan, Tarantola, Saltelli, and Sobol 2000 for a review). Statisticians, and practitioners of experimental design, call V_i the first-order effect of X_i on Y .

3.1 Noncorrelated Factors

The factors X_i are noncorrelated. Assume that V_l and V_r , related to X_l and X_r , are both smaller than V_j , associated with X_j . However, X_l and X_r may be involved in an interaction effect that is not captured by V_l or V_r . How can we establish whether X_j is actually more important than X_l and X_r in lottery setting 1? The interaction effect between X_l and X_r is given by

$$V_{lr} = V[E(Y|X_l, X_r)] - V[E(Y|X_l)] - V[E(Y|X_r)], \quad (8)$$

where $V[E(Y|X_l, X_r)]$ is the joint effect of X_l and X_r and V_{lr} is the pure interaction of X_l and X_r , known as a second-order (or two-way) effect. Analogous formulas can be written for higher-order terms. If there are no interactions, then

$$V = \sum_j V_j \quad (9)$$

and we can establish the relative importance of the factors using the respective V_j values. If the model contains interactions, then

$$V = \sum_i V_i + \sum_{i<j} V_{ij} + \sum_{i<j<m} V_{ijm} + \dots + V_{12\dots k}; \quad (10)$$

that is, the sum of all the first-order and higher-order terms adds up to the total unconditional variance V (Sobol' 1990). V_{ijm} is the third-order interaction among X_i , X_j , and X_m . $V_{12\dots k}$ is the k th-order interaction among all of the factors. It can be easily shown that, even when higher-order terms in (10) are nonzero, the V_i are sufficient to make an informed choice on lottery setting 1, based on the very definition of this measure. If $V_j = V[E(Y|X_j)]$ is greater than V_l and V_r , then V_j is the most important factor, regardless of the value of V_{lr} . For V_{lr} to come into play, one should stipulate a setting in terms of "what couple of factors should be fixed" and so forth.

Conclusion 1. In lottery setting 1, for noncorrelated factors, the V_j 's are the measure to use to make an informed choice, whether or not the factors interact.

3.2 Correlated Factors

When the X_j are correlated, the output variance cannot be decomposed as in (10). Furthermore, the V_j are more expensive to obtain because the computational shortcuts available for noncorrelated inputs (see, e.g., Sobol 1990) are no longer applicable. However, a rather efficient estimation procedure for the first-order terms due to McKay (1995) uses the replicated Latin hypercube sampling (LHS) design (r -LHS). The correlation in the r -LHS sample is induced using the permutation procedure proposed by Iman and Conover (1982).

Leaving aside the computational issues, and going back to our lottery setting 1, we observe that V_j is again the measure to use for making an informed choice, even when the input is correlated. This descends from the definition of the setting, which implies the "fixing" of a single factor when computing the conditional variance. Note that V_j will be sensitive in general to the existence of correlations. Another observation relevant to the case of correlated input is that although in lottery setting 1, the fact that factors are fixed one at a time would normally prevent the detection of interactions, yet in the presence of correlation, fixing one factor also influences the distribution of the others. This may allow the influence of interactions to emerge, depending on the relative patterns of correlation and interactions. As a trivial example, if the model is $Y = x_1 + x_2 + x_3 + a_{23} * x_2 x_3$ and x_1 and x_2 are correlated, then $V(E(Y|x_1))$ will depend on a_{23} as well as on the distribution parameters of x_2 and x_3 . In this sense, we say that interactions may be "carried over" by correlation. This effect is a possibility only when the input is correlated, and is absent when the input is not correlated. As a result, an important consequence of adopting lottery setting 1 is that we accept the risk of remaining ignorant about important features of the model that are the objects of the sensitivity analysis: the presence of interactions. We are hence ready to offer our second conclusion.

Conclusion 2. In Lottery Setting 1, the terms V_j are the measures that guarantee an informed choice in the cases where the factors do not interact and do not correlate, interact but do not correlate (Conclusion 1), and correlate and interact.

4. LOTTERY SETTING 2

This context is relevant to the broad area of risk assessment and management, to which Level E belongs. Here one seeks to achieve a preestablished reduction in the variance of Y .

Lottery Setting 2. We are asked to bet on sets of factors (couples, triplets, and so on). The prize is for obtaining a variance of Y equal or smaller than a given target variance, $V_{tar} < V$, by fixing simultaneously the smallest number of factors.

We introduce first a more compact notation, with the superscript c standing for “closed” within a subset of factors. For instance, $V_{jm}^c = V[E(Y|X_j, X_m)]$ and $V_{npq}^c = V[E(Y|X_n, X_p, X_q)]$.

4.1 Noncorrelated Factors

Assume that the model is additive (i.e., there are no interactions) and that the target reduced variance is $V_{tar}/V = .1$, which corresponds to 90% reduction in the variance of Y . Given that $V = \sum_i V_i$, the factors can be ranked in order of importance using the V_i (i.e., $V_{R1} \geq V_{R2} \geq \dots \geq V_{Rk}$). Then we select the first r factors such that $\sum_{i=1}^r V_{Ri} \geq V - V_{tar}$.

If the model is nonadditive (i.e., there are interactions between factors), then we could use this procedure to select the smallest set of V_{i_1, i_2, \dots, i_s} ’s such that their sum just about exceeds $V - V_{tar}$:

1. Take the factor with the highest V_i . If $V_i > V - V_{tar}$, then end the procedure; otherwise, go to step 2.
2. Evaluate all of the second-order partial variances (i.e., $k(k-1)/2$ terms) and take the pair X_j and X_m with the highest V_{jm}^c (neither X_j nor X_m must necessarily coincide with the X_i considered in step 1). If $V_{jm}^c > V - V_{tar}$, then choose X_j and X_m ; otherwise, go to step 3.
3. Evaluate all of the third-order partial variances (i.e., $k(k-1)(k-2)/6$ terms) and take the triplet X_n, X_p, X_q with the highest V_{npq}^c . If $V_{npq}^c > V - V_{tar}$, then choose X_n, X_p, X_q , and so on with higher-order terms.

The number of terms V_{i_1, i_2, \dots, i_s} required is not fixed a priori, and the procedure might be computationally impracticable even for moderate values of k . We recommend a different strategy for this setting that requires using the total sensitivity index $S_{Ti} = V_{Ti}/V$, where the V_{Ti} for a given factor X_i is the sum of all of the terms in (10) that include the index i . For instance, in a model with three factors, the V_{Ti} are

$$\begin{aligned} V_{T1} &= V_1 + V_{12} + V_{13} + V_{123}, \\ V_{T2} &= V_2 + V_{12} + V_{23} + V_{123}, \\ V_{T3} &= V_3 + V_{13} + V_{23} + V_{123}. \end{aligned} \tag{11}$$

V_{Ti} can also be defined as $E[V(Y|X_{-i})]$, representing the average variance that would remain as long as X_i stays unknown. Note that X_{-i} indicates all of the factors but X_i . The V_{Ti} can be computed without computing the single terms in

(11) (Homma and Saltelli 1996). The strategy that we recommend in the presence of interactions is as follows:

1. Compute the full set of V_i ’s and V_{Ti} ’s.
2. Rank the factors by using the V_{Ti} , obtaining a sequence V_{TR1}, V_{TR2}, \dots such that $V_{TR1} > V_{TR2} > \dots$.
3. Take the factor with the highest total index, that is, X_{R1} . If $V_{R1} > V - V_{tar}$, then end the procedure. Otherwise, go to step 4.
4. Enter the factor with the second-highest total index, that is, X_{R2} . If $V_{R1, R2}^c > V - V_{tar}$, then end the procedure, and so on.

This procedure is cheaper than the previous one, although it could become cumbersome if it did not converge rapidly. An even cheaper alternative could be to reach the threshold $V - V_{tar}$ by using the V_i ’s alone instead of the closed terms. But for highly nonadditive models, this could lead to overshooting, because fixing X_i and X_j removes V_{ij} besides removing V_i and V_j ; fixing X_i, X_j, X_k removes V_{ij}, V_{ik}, V_{jk} , and V_{ijk} besides removing V_i, V_j , and V_k ; and so on.

4.2 Correlated Factors

If the input factors are correlated, then we are not allowed to relate $V_{ir}^c = V[E(Y|X_i, X_r)]$ to the sum of a “purely interaction” term V_{ir} and the individual effects V_i and V_r , as in (8), although we can still compute V_{ir}^c . Furthermore, if either X_i or X_r are correlated with a third factor X_j , then the V_{ir}^c measure will depend on the strength of such correlation as well as on the distribution of X_j .

It should also be recalled that the estimation of the conditional variances of any order is much more expensive computationally in case of correlated input (see Sec. 5.4 for details). For this reason, the number of conditional variances must be kept to a minimum. Our proposed procedure for correlated input is as follows:

1. Rank the factors in order of importance using the V_i and obtain a sequence $V_{R1}, V_{R2}, \dots, V_{Rk}$, where $V_{R1} \geq V_{R2} \geq \dots \geq V_{Rk}$. If $V_{R1} \geq V - V_{tar}$, then X_{R1} is the choice; end the procedure. Otherwise, go to step 2.
2. Compute the second-order term V_{R1j}^c , that is, the reduction of the variance of Y that can be achieved by fixing the pair (X_{R1}, X_j) . X_j is the factor that exhibits the highest “figure of merit” M_j (defined later). If $V_{R1j}^c > V - V_{tar}$, then (X_{R1}, X_j) is the choice; otherwise, go to step 3.
3. Compute the third-order term V_{R1jm}^c , where X_m is selected using the figure of merit. If $V_{R1jm}^c > V - V_{tar}$, then (X_{R1}, X_j, X_m) is the choice, and so on.

The figure of merit M_j is defined as

$$M_j = S(R_j) \left(1 - \max_{i \in u} |c_{ij}| \right) \left(1 + \frac{V_{Tj}^{NC} - V_j^{NC}}{V^{NC}} \right)^2, \tag{12}$$

where $S(R_j) = \sum_{r=R_j}^k \frac{1}{r}$ is a Savage score (Savage 1956). Equation (12) means that the rank R_j of the candidate factor X_j provided at step 1 is first converted into a Savage score and is then penalized or prized depending on its correlations and interactions. Here u is the set of the input factors already

selected, and c_{ij} is the correlation coefficient between X_i and X_j , which is known a priori. A candidate factor for inclusion in u that is correlated with some of the factors already in u must be penalized, because part of its effect on the output variance might have already been offset by fixing the set in u . This penalization should be proportional to the degree of correlation. The resulting score is then prized if X_j interacts with other factors already in u , because this could lead to a higher variance reduction than is implied by its V_j . The extent of the interaction is quantified by $V_{T_j}^{NC} - V_j^{NC}$, where the superscript "NC" means that the measures are computed for the noncorrelated problem. Empirical tests have shown that the figures of merit are effective when the prize for interaction is squared.

5. THE LEVEL E IN SETTINGS 1 AND 2

5.1 Setting 1 With Noncorrelated Factors

The output of interest is the total radiologic dose, $Y(t)$, from (6), calculated at $t = 10^5$ yr in the future. This time point is special because it corresponds to the lowest value for the model coefficient of determination on ranks R_y^{*2} (Fig. 1), where the model shows a strong nonadditive, nonmonotonic behavior.

We express computational costs in terms of model evaluations, that is, the number of times the model must be evaluated. Because Y may be the output of an expensive computer code, it is clear that effective sensitivity analysis methods must strive to keep this number low. The method of Sobol' (Sobol' 1990), improved by Saltelli (2002), is used to estimate all of the V_j , and hence the first-order sensitivity indices $S_j = V_j/V$.

A base sample of 1,024 points is considered. The total cost needed to estimate all the V_j is $1,024 \times (12 + 2) = 14,336$ model evaluations, where two extra samples of 1,024 points each are taken to obtain two estimates of the output mean. (Details on sample size and computational cost have been given in Saltelli 2002.)

The analysis shows that the most important factors are $v^{(1)}$ and $R_I^{(1)}$, with S_j equal to .18 and .09. The sum of all the S_j is .38, meaning that the first-order effects explain 38% of the uncertainty in the total dose, with the remaining portion explained by interactions.

The ideal use for the lottery setting 1 is for the prioritization of research, one of the most common uses of sensitivity analysis. Under the hypothesis that all uncertain factors are susceptible for determination (at the same cost per factor), lottery setting 1 allows identification of the factor most deserving an experimental measurement. If we are in a forecast (prognostic) setting and want to reduce the overall uncertainty in our predicted dose, then $v^{(1)}$ is the factor whose uncertainty should be reduced first. If we are in a diagnostic or calibration setting, then again $v^{(1)}$ is the factor that has the best chance of being calibrated. The total cost of this exercise is 14,336 model runs.

5.2 Setting 1 With Correlated Factors

Let us consider Level E with the input correlation structure given in Table 2. The output of interest is again the total radiologic dose, $Y(t)$, at $t = 10^5$ yr. The pattern of R_y^2 and R_y^{*2} across time is almost identical to that in Figure 1 for the noncorrelated case.

We have estimated all of the V_j using r -LHS. An r -LHS consists of r replicates of the LHS obtained as suitable permutations of all the columns of a base LHS sample of size n . We used $n = 500$ and $r = 20$. The results are listed in Table 3. With respect to the noncorrelated case, $v^{(1)}$ has lost its leadership due to the high correlation with T and $v^{(2)}$, which are less influential. This result shows that the omission of correlations in the original specification of the Level E exercise was legitimate as far as uncertainty analysis is concerned (the output dose does not change much), but was an oversimplification as far as the identification of the influential factors is concerned.

Note that for the noncorrelated case of Level E, $\sum_j V_j = 2 \times 10^{-15}$ and $V = 5.3 \times 10^{-15}$, the difference being attributable




Figure 1. Model Coefficient of Determination for the Level E Case Study as a Function of Time (yr). Circles refer to the values of R_y^2 ; diamonds, to the values of R_y^{*2} (i.e., obtained using the ranks of the model output values). The output variable is the total annual dose, Y . The poor performance of the regression model is highlighted by the very small value of R_y^2 at every time point. At $t = 10^5$ yr, R_y^{*2} also is very low, indicating that even rank-based statistics are inappropriate for the purpose of sensitivity analysis. R_y^2 and R_y^{*2} have been estimated using the sequence of output values obtained by running the Level E model on a Latin hypercube sample of size 1,000 generated over the space of the inputs.

Table 3. Lottery Setting 1: The V_j and the Ratios $S_j = V_j/V$ for the Correlated Case

Factor	$V_j \cdot 10^{17}$	$S_j = V_j/V$
W	91.0	.08
$v^{(1)}$	38.4	.03
$R_I^{(1)}$	11.0	.01
T	8.7	.01
$R_C^{(2)}$	6.8	.01
$v^{(2)}$	6.6	.01
$I^{(2)}$	6.4	.01
$I^{(1)}$	6.1	.01
$R_C^{(1)}$	5.9	.00
$R_I^{(2)}$	2.1	0
k_C	1.4	0
k_I	1.0	0

NOTE: The factors are ranked in decreasing order of importance with respect to the S_j .

to interactions. For the correlated case $\sum_j V_j = 1.85 \times 10^{-15}$ and $V = 1.18 \times 10^{-14}$. When input is correlated, we cannot conclude that the missing fraction of variance is that due to interaction, because (10) does not hold. The fraction due to interaction could in fact be higher than the difference $V - \sum_j V_j$. The total cost of this exercise is 10,000 model runs.

5.3 Setting 2 With Noncorrelated Factors

We now apply the procedure of Section 4.1 to Level E with noncorrelated input. We assume that the targeted variance reduction in the model output is 40%. The technique proposed by Saltelli (2002) enables us to estimate all of the V_j and the V_{T_j} (step 1 of the procedure) for the 12 factors at the cost of 14,336 model runs, as in Section 5.1. The model is strongly nonadditive given that, as mentioned in Section 5.1, $\sum_j V_j^{NC}/V = .38$. Sizeable variance reductions can be achieved only by fixing more factors simultaneously. The top-ranked factor is $v^{(1)}$ with the highest $V_{T_j}^{NC}$ (step 2), but $V_{v^{(1)}}^{NC}/V = .18$ is below the target imposed (step 3). W is the factor with the second-highest $V_{T_j}^{NC}$; we estimate $V_{v^{(1)}W}^c$ with the method of Sobol' at the cost of 1,024 model evaluations (step 4). The target variance reduction is still not reached (28%). We need to enter the third-ranked factor, $l^{(1)}$, and estimate $V_{v^{(1)}l^{(1)}W}^c$ using the method of Sobol' with an other 1,024 simulations. The variance reduction obtained by simultaneously fixing $v^{(1)}$, W , and $l^{(1)}$ is 51%, which enables us to meet the target. This result is remarkable and novel. It shows that we can achieve a considerable overall variance reduction by judiciously fixing a set of interacting factors (a triplet in this case). To a certain extent, we can generalize these results to problems of mass transfer with chemical reaction. We know that factors that are downstream in the modeling chain and that influence the output linearly (W) tend to interact with those that govern the transit time, $v^{(1)}$ and $l^{(1)}$. The total cost of this exercise is $14,336 + 1,024 + 1,024 = 16,384$ model runs.

5.4 Setting 2 With Correlated Factors

We now apply the procedure of Section 4.2 to Level E with correlated input, with the same target variance reduction (40%). The ranking provided by the V_j (step 1) yields W as the top-ranked factor with $S_i = 8\%$ (see Table 3). The cost of estimating the V_j is 10,000 runs, as explained in Section 5.2. We compute the figures of merit M_j as from (12) (Table 4). This requires other 14,336 runs, as explained in Section 5.3. The largest M_j is that for $v^{(1)}$. According to step 2, we compute $V_{Wv^{(1)}}^c$. To obtain this estimate, we must compute the inner conditional expectation $E(Y|W, v^{(1)})$ through a numerical integration over a base sample of, say, $N_b = 100$ points. The same conditional expectation has to be evaluated over a set of different values of the pair ($W, v^{(1)}$) (say, $N_r = 100$), to compute the variance of the conditional expectations. Therefore, this estimate requires $N = N_b \times N_r = 10,000$ model simulations. The target is not reached, because a variance reduction of 32% is obtained by fixing the pair ($W, v^{(1)}$). The factor $R_l^{(1)}$ (with $M_j = 2.43$) is included in the analysis (step 3), and the term $V_{Wv^{(1)}R_l^{(1)}}^c$ is estimated (requiring another 10,000 runs); the target is reached because the variance reduction rises to about 48%. (The variance reduction

Table 4. Lottery Setting 2: Values of the Ratios V_j^{NC}/V and $V_{T_j}^{NC}/V$ (for the noncorrelated set) and the Figures of Merit M_j for Use in Our Recommended Procedure

Factor	V_j^{NC}/V	$V_{T_j}^{NC}/V$	M_j
W	.04	.55	—
$v^{(1)}$.18	.87	6.01
$R_l^{(1)}$.09	.32	2.43
T	0	0	1.27
$l^{(1)}$.03	.50	1.10
$R_C^{(2)}$	0	0	1.02
$v^{(2)}$.04	.07	.87
$l^{(2)}$	0	.02	.76
$R_C^{(1)}$	0	0	.39
$R_l^{(2)}$	0	0	.27
k_C	0	0	.17
k_l	0	0	.08

NOTE: The factors are ranked in order of decreasing importance with respect to the values of M_j .

grows rapidly if we are able to fix the "right" factors.) The result of the correlation between factors, with respect to the uncorrelated case is that $R_l^{(1)}$ is now a better partner for W and $v^{(1)}$ for obtaining the maximum output variance reduction. If $l^{(1)}$ instead of $R_l^{(1)}$ were fixed, then the output variance reduction would be less, that is, 43%. The total cost of this exercise is $10,000 + 14,336 + 10,000 + 10,000 = 44,336$ model runs.

6. CONCLUSIONS

The motivation of this article is to devise a strategy for sensitivity analysis that could work even in the simultaneous presence of correlated input and nonadditive models. To establish objective criteria for the evaluation of the proposed strategy, we have tried to define in a rigorous fashion the questions posed by the sensitivity analysis. The literature shows that many different approaches and framing assumptions result in different definitions of importance and sensitivity. Without a rigorous problem setting, alternative definitions of importance are equally plausible (see OECD 1993).

The two lottery settings proposed in this article cover important areas of application for sensitivity analysis. The first setting can be used for priority setting, to identify which factor has the greatest potential for output uncertainty reduction through additional research. The setting also applies to calibration analysis; it would help to design an experiment in such a way that the factors to be measured, in the presence of other nonreducible uncertainties, are the most important in lottery setting 1. For this setting, the V_j 's represent the proper measure to use whatever the correlation and interaction structure of the model.

The second setting looks at the compounded effects of factors in a model and could represent the setting in which "risk" is involved. Extreme events, those of interest to risk analysts, often occur when interacting factors assume their most unfavorable values concurrently. Lottery setting 2 aims to identify the smallest number of factors to be fixed to obtain the maximum reduction in the variance of the output. Control theory is another area in which this setting might be of use.

For the two settings, we have proposed procedures to fight the curse of dimensionality, for example, the fact that the number of conditional variances in a model grows exponentially

with problem dimensionality. It is clear that importance assessments are more cumbersome for correlated inputs than for noncorrelated inputs, as many more evaluations of the model are needed. In the procedure suggested in Section 4.2, the number of conditional variances to be estimated remains on the order of k . Better (although much more expensive) results can be obtained by computing all possible conditional variances, $O(2^k)$, or using the procedure proposed by McKay, Morrison, and Upton (1999), $O(k(k-1)/2)$.

The high computational cost of quantifying importance of input factors makes the approach described in this article unfeasible for computationally expensive models (1 hour or more of CPU time per simulation, say). The present approach is instead reasonable for models in the seconds-to-minute range, which represent the vast majority of models. The test case addressed in this work offered two main practical advantages. On the one hand, it is very well known to practitioners in risk assessment and sensitivity analysis; on the other hand, it is representative of a large class of environmental or risk analysis models, that deal substantially with a problem of mass transfer with chemical reaction. In our experience, for instance, the pattern of interaction displayed by Level E is very representative of situations found in hydrology, air pollution, and nuclear safety studies.

[Received June 2000. Revised April 2002.]

REFERENCES

- Chan, K., Tarantola, S., Saltelli, A., and Sobol', I. M. (2000), "Variance-Based Methods," in *Sensitivity Analysis*, eds. A. Saltelli, K. Chan, M. Scott, New York: Wiley, pp. 167–197.
- CPC, Computer Physics Communications, (1999). Special Issue on Sensitivity Analysis, 117(1–2), 1–188.
- Crank, J. (1975). *The Mathematics of Diffusion*, Oxford, UK: Oxford University Press.
- Draper, D., Pereira, A., Prado, P., Saltelli, A., Cheal, R., Eguilior, S., Mendes, B., and Tarantola, S. (1999), "Scenario and Parametric Uncertainty in GESAMAC: A Methodologic Study in Nuclear Waste Disposal Risk Assessment," *Computer Physics Communications*, 117, 142–155.
- Draper, D., Saltelli, A., Tarantola, S., and Prado, P. (2000), "Scenario and Parametric Sensitivity and Uncertainty Analyses in Nuclear Waste Disposal Risk Assessment: The Case of GESAMAC," in *Sensitivity Analysis*, eds. A. Saltelli, K. Chan, and M. Scott, New York: pp. 275–292.
- Homma T., and Saltelli, A. (1996), "Importance Measures in Global Sensitivity Analysis of Model Output," *Reliability Engineering and System Safety*, 52, 1–17.
- Hora, S. C., and Iman, R. L. (1989), "Expert Opinion in Risk Analysis: The NUREG-1150 Methodology," *Nuclear Science and Engineering*, 102, 323–331.
- Iman, R. L., and Conover, W. J. (1982), "A Distribution Free Approach to Inducing Rank Correlation Among Input Variables," *Communications in Statistics, Part B—Simulation and Computation*, B11, 311–334.
- Ishigami, T., and Homma, T. (1990), "An Importance Quantification Technique in Uncertainty Analysis for Computer Models," in Proceedings of the ISUMA '90, First International Symposium on Uncertainty Modelling and Analysis.
- JMCDA, *Journal of Multicriteria Decision Analysis*, (1999), Special Issue on Sensitivity Analysis, D. Rios Insua (Ed.), Vol. 8.
- JSCS, *Journal of Statistical Computation and Simulation*, (1997), Special Issue on Sensitivity Analysis, 57(1–4).
- McKay, M. D. (1995), "Evaluating Prediction Uncertainty," Report NUREG/CR-6311, U.S. Nuclear Regulatory Commission and Los Alamos National Laboratory.
- McKay, M. D., Morrison, J. D., and Upton, S. C. (1999), "Evaluating Prediction Uncertainty in Simulation Models," *Computer Physics Communications*, 117, 44–51.
- OECD (1989), "OECD/NEA PSAC User Group, PSACOIN Level E Intercomparison. An International Code Intercomparison Exercise on a Hypothetical Safety Assessment Case Study for Radioactive Waste Disposal Systems," technical report, OECD—NEA, Paris.
- OECD (1993), "OECD/NEA PSAG User Group, PSACOIN Level S Intercomparison. An International Code Intercomparison Exercise on a Hypothetical Safety Assessment Case Study for Radioactive Waste Disposal Systems," technical report, OECD—NEA, Paris.
- Prado, P., Homma, T., and Saltelli, A. (1991), "Radionuclide Migration in the Geosphere. A 1D Advective and Dispersive Transport Module for Use in Probabilistic System Assessment Codes," *Radioactive Waste Management and Nuclear Fuel Cycle*, 16, 49–68.
- RESS, Reliability Engineering and System Safety, (1997), Special Issue on Sensitivity Analysis, 57(1).
- Risk Newsletter, (1987), Defining "Risk," Editorial note, *Risk Newsletter* 7(3), September 1987, 5.
- Robinson, P. C., and Hodgkinson, D. P. (1987), "Exact Solutions for Radionuclide Transport in the Presence of Parameter Uncertainty," *Radioactive Waste Management and the Nuclear Fuel Cycle*, 8, 283–311.
- Sacks J., Schiller, S., and Welch, W. (1989), "Designs for Computer Experiments," *Technometrics*, 31, 41–47.
- Sacks, J., Welch, W. J., Mitchell, T. J., and Wynn, H. P. (1989b), "Design and Analysis of Computer Experiments," *Statistical Science*, 4, 409–435.
- Saltelli, A. (2002), "Making Best Use of Model Evaluations to Compute Sensitivity Indices," *Computer Physics Communications*, 145, 280–297.
- Saltelli A., Andres, T. H., and Homma, T. (1993), "Some New Techniques in Sensitivity Analysis of Model Output," *Computational Statistics and Data Analysis*, 15, 211–238.
- Saltelli, A., and Bolado, R. (1998), "An Alternative Way to Compute the Fourier Amplitude Sensitivity Test (FAST)," *Computational Statistics and Data Analysis*, 26, 445–460.
- Saltelli A., Chan, K., and Scott, M. (Eds.) (2000), *Sensitivity Analysis*, New York: Wiley.
- Saltelli, A., and Homma, T. (1992), "Sensitivity Analysis for Model Output. Performance of Black-Box Techniques on Three International Benchmark Exercises," *Computational Statistics and Data Analysis*, 13, 73–94.
- Saltelli, A., and Sobol', I. M. (1995), "About the Use of Rank Transformation in Sensitivity Analysis of Model Output," *Reliability Engineering and System Safety*, 50, 225–239.
- Saltelli, A., Tarantola, S., and Campolongo, F. (2000), "Sensitivity Analysis as an Ingredient of Modelling," *Statistical Science*, 15, 1–20.
- Saltelli, A., Tarantola, S., and Chan, K. (1999), "A Quantitative, Model Independent Method for Global Sensitivity Analysis of Model Output," *Technometrics*, 41, 39–56.
- Savage, I. R. (1956), "Contributions to the Theory of Rank Order Statistics—The Two-Sample Case," *The Annals of Mathematical Statistics*, 27, 590–615.
- Sobol', I. M. (1990), "Sensitivity Estimates for Nonlinear Mathematical Models," *Matematicheskoe Modelirovanie*, 2, 112–118, translation (1993), "Sensitivity Analysis for Nonlinear Mathematical Models," *Mathematical Modelling & Computational Experiment*, 1, 407–414.
- Welch, W. J., Buck, R. J., Sacks, J., Wynn, H. P., Mitchell, T. J., and Morris, M. D. (1992), "Screening, Predicting, and Computer Experiments," *Technometrics*, 34, 15–47.