



Methodology for comparative model assessment and validation

By Andrea Saltelli, at the Faculty of Management and Economics of the Gdańsk University of Technology (Politechnika Gdańska), October 28 2026 – 28 października 2026, godz. 13:15–15:00, sala 103/104 WZiE



Based on Chapter 5 of Equations of Power: Models, Authority, and the Politics of Knowledge (Anthem Press, 2027)



“Ce qui est simple est toujours faux. Ce qui ne
l’est pas est inutilisable.”

(“If it’s simple, it’s always false. If it’s not, it’s unusable.”)

— Paul Valéry, Œuvres II (1942)

Sensitivity analysis and validation

“But the real strength of the models, in my mind at least, were in *sensitivity analysis* where one could examine the response of the model to parameters or structures that were not known with precision (i.e., sensitivity analysis), and in the examination of the behavior of the model components relative to that of the real system in question (i.e., *validation*). By undertaking sensitivity analysis and validation, a great deal can be learned about the real system, including what you do not know.” (*Italics* by Hall)



Charles A. S. Hall, ecologist

The map for today: technical versus normative quality





Figure 5.1, Equations of Power: Models, Authority, and the Politics of Knowledge (Anthem Press, 2027).


Part I – Global sensitivity analysis as a severe test



Karl Popper
(1902–1994)

Models that take no risks

Popper: The empirical content of a theory is measured by the risks it takes — the observations that could prove it wrong (severe tests)




- Climate models projecting the cost of CO₂ to 2300.
- Nuclear waste safety models covering a million years.
- Digital twins simulating ecosystem dynamics with 50-year horizons

These models take no risks. No severe tests here

Uncertainty analysis: the study of the uncertainty in model output




The cascade of uncertainty



Source: <https://www.climate-lab-book.ac.uk/2014/cascade-of-uncertainty/>

Sensitivity analysis: the study of the relative importance of different input factors on the model output



Why global? Why don't
we simply move one
factor At a Time (OAT)


A geometric proof



Contents lists available at ScienceDirect

Environmental Modelling & Software

journal homepage: www.elsevier.com/locate/envsoft




How to avoid a **perfunctory** sensitivity analysis

Andrea Saltelli*, Paola Annoni

Joint Research Center, Institute for the Protection and Security of the Citizen, via E.Fermi, 2749, Ispra VA 21027, Italy


OAT in 2 dimensions



Area circle
/ area
square =?

$\sim 3/4$

OAT in 3 dimensions



Volume sphere /
volume cube =?


$\sim 1/2$

OAT in 10 dimensions; Volume
hypersphere / volume ten dimensional
hypercube =?

~ 0.0025



OAT in k dimensions



OAT does not capture interactions



➔ The resulting analysis is non conservative



How would you test the scaffolding?

How coupled ladders are shaken in most of available literature




How to shake coupled ladders






Plotting the output as a function of two different input factors


Which factor is more important?



Each pink point is $\sim E_{\mathbf{x}_{\sim i}}(Y|X_i)$



Which factor
has the highest
 $V_{X_i} \left(E_{\mathbf{x}_{\sim i}} (Y | X_i) \right) ?$



$$S_i = \frac{V_{X_i} \left(E_{\mathbf{X}_{\sim i}} (Y | X_i) \right)}{V(Y)}$$

The partial variance divided by the total variance is the so-called sensitivity index of the first order

Total effect term

$$T_i = \frac{V_{X_i} \left(E_{\mathbf{X}_{\sim i}}(Y | X_i) \right)}{V(Y)}$$


Homma, Toshimitsu, and Andrea Saltelli. 1996. 'Importance Measures in Global Sensitivity Analysis of Nonlinear Models'. *Reliability Engineering & System Safety* 52 (1): 1–17. [https://doi.org/10.1016/0951-8320\(96\)00002-6](https://doi.org/10.1016/0951-8320(96)00002-6).

Given a 3-factor model $f(X_1, X_2, X_3)$

Where the variance decomposition would

$$\text{read } 1 = S_1 + S_2 + S_3 + S_{12} + S_{13} + S_{23} + S_{123}$$

We compute



$$T_1 = S_1 + S_{12} + S_{13} + S_{123}$$

$$T_2 = S_2 + S_{12} + S_{23} + S_{123}$$

$$T_3 = S_3 + S_{13} + S_{23} + S_{123}$$

“Sensitivity analyses would help” (Leamer, 1985)

JOURNAL ARTICLE

Sensitivity Analyses Would Help

Edward E. Leamer



The American Economic Review

Vol. 75, No. 3 (Jun., 1985), pp. 308-313 (6 pages)

Published By: American Economic Association

<https://www.jstor.org/stable/1814801>

GSA as a severe test

Does the conclusion survive a test designed to refute it?

When the inference dissolves under the exploration of its own input space, the model's lack of empirical content is exposed.

Edward E. Leamer, 2010

“One reason these methods are rarely used is their honesty seems destructive; or, to put it another way, a fanatical commitment to fanciful formal models is often needed to create the appearance of progress.”

GROPING in the DARK


THE FIRST DECADE OF
GLOBAL MODELLING

Donella Meadows
John Richardson
Gerhart Bruckmann

“We call a parameter sensitive **not** when a change in it changes the value of an output number but when it changes the policy conclusions we would draw from the model” (p. 132)



Giandomenico Majone



- Data
- Analyst
- Model
- Objective

“Are the results from a particular model more sensitive to changes in the model and the methods used to estimate its parameters, or to changes in the data?”

Which cause should drive the result?

Material cause = the data

The result should depend on it

Final cause = the objective

The result should depend on it

Formal cause = the model


The result should depend on it least

Efficient cause = the analyst

The result should depend on it least

Models should be sensitive to the data and the objective, and insensitive to the model and the analyst using it.

The Song Jian / one-child policy 1980's model was found to be near-insensitive to inputs ...



Download This Paper

Open PDF in Browser

What Demographic Prediction Can and Cannot Achieve

14 Pages • Posted: 19 May 2026

[Samuele Lo Piano](#)

Gdańsk University of Technology - Faculty of Management and Economics

[Marta Kuc Czarnecka](#)

Gdańsk University of Technology

[Roger Pielke](#)


University of Colorado at Boulder

[Andrea Saltelli](#)

UPF Barcelona School of Management


Date Written: May 15, 2026

... and projections **today**
in use are dramatically
model-dependent



Current Models Underestimate Future Irrigated Areas

- How much land will need to be irrigated by the year 2050?
- Here the dashed lines represent deterministic model predictions from different models and datasets (from FAO & others organizations);
- An uncertainty analysis (grey histogram) reveals that the models are non-conservative: the need might be much larger



Citation:

Puy, A., Lo Piano, S., & Saltelli, A. (2020). Current models underestimate future irrigated areas. *Geophysical Research Letters*, 47, e2020GL087360. <https://doi.org/10.1029/2020GL087360>


Part II – From one model to many

Robustness analysis (Houkes et al., 2024)

parameter RA → dependence on model parameters

structural RA → dependence on structural assumption

representational RA → model philosophy, e.g., moving from a math / differential equations to agent-based modelling



Wybo Houkes, Dunja Šešelja, and Krist Vaesen. Robustness Analysis. In Tarja Knuuttila, Natalia Carrillo, and Rami Koskinen, editors, The Routledge Handbook of Philosophy of Scientific Modeling, pages 195–207. Routledge, Abingdon, Oxon ; New York, NY, 2024.

Robust theorems: “the intersection of independent lies”

“Therefore, we attempt to treat the same problem with several alternative models each with different simplifications but with a common biological assumption. Then, if these models, despite their different assumptions, lead to similar results we have what we can call a robust theorem which is relatively free of the details of the model. Hence our truth is the intersection of independent lies.” (Levins, 1966, p. 423)

Orzack & Sober (1993)

Critical: Robustness is a non-empirical form of confirmation


Weisberg (2006)

Supportive: A four-step procedure linking common structures across models to robust properties

What GSA adds

The common structure must also be shown to be influential

Perturbed parameter (or physics) ensemble: same as uncertainty quantification and sensitivity analysis when done by physicists in climate modelling



John W. Rostron, David M. H. Sexton, Carol F. McSweeney, Kuniko Yamazaki, Timothy Andrews, Kalli Furtado, Mark A. Ringer, and Yoko Tsushima. The impact of performance filtering on climate feedbacks in a perturbed parameter ensemble. *Climate Dynamics*, 55(3):521–551, August 2020. doi:10.1007/s00382-020-05281-8

Butterflies and hawkmoths

The butterfly effect

Sensitive dependence on initial conditions (SDIC)

Can be explored by propagating uncertainty

The hawkmoth effect


Structural model error (SME)

Imposes often terminal limits on predictive power (Frigg et al., 2014)

Yet structural uncertainty **can itself be propagated** — e.g. by sampling among alternative formulations of the same physical law (evapotranspiration).

Via negativa (N.N. Taleb's books): such analyses establish that a model cannot support the inference asked of it — and may cure the modeller of hubris.

Sample modelling choices via 'triggers'. Global sensitivity analysis becomes thus a modelling of the modelling process = multiverse analysis



Example of using triggers: an integer in $[1-5]$ is used to select one out of 5 possible evapotranspiration formulae at each model run

nature communications

[Explore content](#) ▾ [About the journal](#) ▾ [Publish with us](#) ▾

[nature](#) > [nature communications](#) > [comment](#) > article

[Comment](#) | [Open access](#) | Published: 08 June 2022


The delusive accuracy of global irrigation water withdrawal estimates

[Arnald Puy](#) , [Razi Sheikholeslami](#), [Hoshin V. Gupta](#), [Jim W. Hall](#), [Bruce Lankford](#), [Samuele Lo Piano](#), [Jonas Meier](#), [Florian Pappenberger](#), [Amilcare Porporato](#), [Giulia Vico](#) & [Andrea Saltelli](#)

[Nature Communications](#) **13**, Article number: 3183 (2022) | [Cite this article](#)

10k [Accesses](#) | **79** [Citations](#) | **93** [Altmetric](#) | [Metrics](#)

Trigger=an integer variable, whose value is determined via 'Russian roulette' in Monte Carlo jargon



Russian roulette as depicted in the 1925 film *The Night Club*. Source: Wikipedia Commons

The garden of forking paths: Why multiple comparisons can be a problem, even when there is no “fishing expedition” or “p-hacking” and the research hypothesis was posited ahead of time*

Andrew Gelman[†] and Eric Loken[‡]

14 Nov 2013

PNAS

RESEARCH ARTICLE

SOCIAL SCIENCES

 OPEN

Observing many researchers using the same data and hypothesis reveals a hidden universe of uncertainty

Edited by Douglas Massey, Princeton University, Princeton, NJ; received March 6, 2022; accepted August 22, 2022

Observing many researchers using the same data and hypothesis reveals a hidden universe of uncertainty

Edited by Douglas Massey, Princeton University, Princeton, NJ; received March 6, 2022; accepted August 22, 2022

The political orientation of the analysts was found **not** to explain much of the variance



The modelling variables themselves were found to explain at most a few per cent of the variance of the results ... Where **does the variance come from?**



<https://www.istockphoto.com/>

Our sensitivity analysis explains where the missing variance went: interactions

Sobol' indices S_i T_i



Many models fit the same data $\cdots \rightarrow$ and tell different stories

Equifinality

Hydrology (Beven, 1993)

The Rashomon effect

Machine learning (Breiman, 2001)


The importance of the same variable may vary across models

→ a sensitivity analysis that keeps the model fixed may be incomplete

→ hence: modelling of the modelling process

Part III – Validation of what? The limits of GSA

The paradox of global sensitivity analysis



“The first act of selection – a sort of proto-sensitivity analysis – occurs in the mind of the modeller, who chooses which aspects of a system are ‘relevant’.”

GSA then chases which of those aspects drive the results — but the second selection is subordinate to the first.



It tests the fabric of the model, rather than the model against the world it claims to represent.



Mary S. Morgan



Margaret Morrison




Models as mediators between theories and the world.

World in the model, model in the world





<<It is important, however, to recognize that the sensitivity of the parameter in the equation is what is being determined, not the sensitivity of the parameter in nature>>

Orrin H. Pilkey



Useless Arithmetic: Why Environmental Scientists Can't Predict the Future
by Orrin H. Pilkey and Linda Pilkey-Jarvis, Columbia University Press, 2009.



The confidence of the stakeholders in TSPA was not helped when evidence was produced which could lead to an upward revision of 4 orders of magnitude of this parameter (the ^{36}Cl story)

In the case of TSPA (Yucca mountain) a range of 0.02 to 1 millimetre per year was used for percolation of flux rate.

→... SA useless if it is instead ~ 3,000 millimetres per year.



Where does the rabbit come from?

A critique: “Monte Carlo doesn’t tell you anything that wasn’t already baked into the simulation. The distribution of the output comes from assumptions in the input (modulo bugs): a probability model for the parameters in the simulation. It comes from what you program the computer to do. Monte Carlo reveals the consequences of your assumptions about randomness. The rabbit goes into the hat when you build the probability model and write the software. The rabbit does not come out of the hat without having gone into the hat first.”
(Philip B. Stark)

A question: How liberal is liberal enough?

Give a factor the widest plausible range and find it non-influential: you have learned something of value. Drop plausibility and let uncertainty explode: what did you learn?

The quality of an analysis $\leftrightarrow$ Its input

From: Uncertainty and Quality in Science for Policy by Silvio Funtowicz and Jerry Ravetz, Springer 1990.

Funtowicz & Ravetz's GIGO (Garbage In, Garbage Out) Science – or pseudo-science – “where uncertainties in inputs must be suppressed least outputs become indeterminate”

Leamer, Edward E. 1985. ‘Sensitivity Analyses Would Help’. The American Economic Review 75 (3): 308–13.

Leamer's ‘Conclusions are judged to be sturdy only if the neighborhood of assumptions is wide enough to be credible and the corresponding interval of inferences is narrow enough to be useful’.

NEVER vary all factors by the same amount
Be it 5%, 10%, or 20%



New WHO estimates: Up to 190 000 people could die of COVID-19 in Africa if not controlled

07 May 2020

Brazzaville – Eighty-three thousand to 190 000 people in Africa could die of COVID-19 and 29 million to 44 million could get infected in the first year of the pandemic if containment measures fail, a new study by the World Health Organization (WHO) Regional Office for Africa finds. The research, which is based on prediction modelling, looks at 47 countries in the




Speculative scenario in which ten uncertain input probabilities are varied by an arbitrary 10% — as if they were truly equally uncertain — with no theoretical or empirical basis for such a choice



“... inferences from mathematical models of phenomena to real physical applications must also be demonstrated to be approximately correct when the assumptions of the model are only approximately true” [1]

Pierre Duhem (1861–1916)
and his ‘Principle of
stability’ ~1906



[1] See: S. C. Fletcher, ‘The Principle of Stability’, *Philosopher’s Imprint*, vol. 20, no. 3, 2020, Accessed: Sep. 08, 2024. [Online].

Part IV – Beyond the formal system: sensitivity auditing and models in competition

From sensitivity analysis to sensitivity auditing

Uncertainty and sensitivity are also, and perhaps eminently, political.

Sensitivity auditing espouses post-normal science and its NUSAP notation: a quantity together with an account of its quality.




Familiar: the number, its units, its range

Judgement of confidence; scored evaluation of the process

What NUSAP adds is not precision but candour.

1. Check against rhetorical use of mathematical modelling
2. Adopt an 'assumption hunting' attitude
3. Detect garbage in, garbage out (GIGO)
4. Find sensitive assumptions before they find you
5. Aim for transparency
6. Do the right sums
7. Focus the analysis on the key question answered by the model, exploring the entire space of the assumptions holistically



Verbatim from

Int. J. Foresight and Innovation Policy, Vol. 9, Nos. 2/3/4, 2013

213


**What do I make of your latinorum? Sensitivity
auditing of mathematical modelling**

Genuine or bogus? A validation checklist

“... [there are] certain criteria by which the mathematician judges whether an “application” of mathematics is genuine or bogus.”

- Does the depth of the real-world problem justify the complexity of the mathematical model?
- Are any genuine mathematical reasonings or non-trivial calculations carried out that require the resources of the mathematical model being proposed?
- Are the coefficients or parameters in the equations capable of being determined in a meaningful and reasonably accurate way?
- Are the conclusions capable of being tested against real-world data?
- Do any non-obvious practical conclusions follow from the analysis?

Eminently sensible — if only one could identify the subject implementing them.



Quantitative story telling: testing the story a model tells

Feasibility

Compatibility with external constraints
e.g. biophysical limits of a resource

Viability

Compatibility with internal constraints
e.g. existing regulatory regimes or laws

Desirability

Compatibility with normative values adopted in the given society
e.g. would citizens willingly submit to the policy?

If the story told by a model is a policy prescription, its robustness is tested against all three — ideally in a participatory setting.

Models in competition: comparative assessment as an institution

Countermodelling

Different modelling teams mobilised by different actors (Zayandeh-Rood basin, Iran)

Adversarial collaboration

Proposed in medicine as a complement to consensus-building

Red teams

Their absence from the UK COVID decision room was lamented by statisticians

Hot tubbing

Concurrent expert evidence in Australian courts: experts in the witness box together

“These instances are instructive precisely because they are exceptional. The default condition remains one where models operate without meaningful challenge.”

None is sufficient on its own

Sensitivity analysis

can reveal what drives a model's output but cannot question the model's purpose.

Sensitivity auditing

can question the purpose but depends on the existence of critical communities willing to do the questioning.

Statactivism and countermodelling

can mobilise such communities but face the asymmetries of power, resources, and access that characterise the present political economy of science.

Fight a number with a number: statactivism

Statistical judo

Gaming the metrics, using Goodhart's law strategically

Denouncing unfair indicators

e.g. the middle-class bias of the French consumer price index

Brand-new measures

A new index that better defends the purchasing power of the poor

Making the invisible visible

Areas of invisibility or exclusion that escape statistical offices




Hall was right: the main reason to build a model may well be to play a sensitivity analysis on it



The question is whether this conflicts with the political economy of mathematical modelling

Decades of scarce traction for opening the black box of models: a debt we now pay with AI

Dona Meadows lamented the scarce traction already in 1985



The End




Backup slides

What would help


A model should not be used for risk analysis unless:

- Its **sensitivity** to key assumptions has been explored Global sensitivity analysis
- Its structural and **framing** choices have been made explicit and tested Modelling of the modelling process
- **Affected parties** have had the opportunity to identify which assumptions matter to them Sensitivity auditing





Ignorantia and via negativa

“Via negativa: the principle that we know what is wrong with more clarity than what is right, and that knowledge grows by subtraction. Also, it is easier to know that something is wrong than to find the fix. Actions that remove are more robust than those that add because addition may have unseen, complicated feedback loops”





Variance-based methods




Plotting the output as a function of two different input factors:

The model has been run 1,000 times sampling all uncertain factors; two of them are shown here, x_i (left) and x_j (right). The output variable has first been sorted by increasing x_i then by increasing x_j .



Same point


The points are always the same, only sorted differently



~1,000 blue points

Divide them in 20 bins of ~ 50 points

Compute the bin's average (pink dots)



$$S_i = \frac{V_{X_i} \left(E_{\mathbf{X}_{\sim i}} (Y | X_i) \right)}{V(Y)}$$

The partial variance divided by the total variance is the so-called sensitivity index of the first order

There are terms which capture two-way, three way, $\cdots$ interactions among variables

All these terms are linked by a formula

Variance decomposition (ANOVA)

$$V(Y) =$$

$$\sum_i V_i + \sum_{i,j>i} V_{ij} + \dots + V_{123\dots k}$$

Variance decomposition (ANOVA)

The total variance can be decomposed into main effects and interaction effects up to the order k , the dimensionality of the problem (only for independent factors)

In fact, interaction terms are awkward to handle: just the second-order terms for a model with k factors are as many as $k(k-1)/2 \dots$

(10 factors=45 second order terms)

How about a single ‘importance’ term for all effects?

In fact, such terms exist and can be computed easily, without knowledge of the individual interaction terms

Before moving to the total index, let us finish with the first-order one

$$S_i = \frac{V_{X_i} \left(E_{\mathbf{X}_{\sim i}} (Y | X_i) \right)}{V(Y)}$$

Known as:

- Pearson's correlation ratio (Pearson)
- First order effect (Sobol' & disciples)
- Top marginal variance (Jansen)

$$S_i = \frac{V_{X_i} \left(E_{\mathbf{X}_{\sim i}}(Y | X_i) \right)}{V(Y)}$$

We need to understand what the numerator is in plain English

$$V_{X_i} \left(E_{\mathbf{X}_{\sim i}}(Y | X_i) \right)$$

$$V_{X_i} \left(E_{\mathbf{x}_{\sim i}} (Y | X_i) \right)$$

Top marginal variance=
 = the expected reduction in variance
 than would be achieved if factor X_i could
 be fixed.

Why?

Because:

$$V_{X_i} \left(E_{\mathbf{X}_{\sim i}} (Y | X_i) \right) + \\ + E_{X_i} \left(V_{\mathbf{X}_{\sim i}} (Y | X_i) \right) = V(Y)$$

One of the exercises in the book: prove this
using $V(\bullet) = E(\bullet)^2 - E^2(\bullet)$

Because:

= then this is the reduction in
variance if X_i could be fixed

$$V_{X_i} \left(E_{\mathbf{X}_{\sim i}} (Y | X_i) \right) +$$

$$+ E_{X_i} \left(V_{\mathbf{X}_{\sim i}} (Y | X_i) \right) = V(Y)$$


(By inspection) this is what variance would be
left (on average) if X_i could be fixed

For additive systems one can decompose the total variance as a sum of first order effects

$$\sum_i V_{X_i} \left(E_{\mathbf{X}_{\sim i}} (Y | X_i) \right) \approx V(Y)$$

How to get from first order to total order

From

$$V_{X_i} \left(E_{\mathbf{X}_{\sim i}} (Y | X_i) \right) \quad \text{Main effect of factor } X_i$$

replacing X_i with $\mathbf{X}_{\sim i}$

Gives the main effect of
non- X_i

$$V_{\mathbf{X}_{\sim i}} \left(E_{X_i} (Y | \mathbf{X}_{\sim i}) \right)$$

But again:

$$V_{\mathbf{X}_{\sim i}} \left(E_{X_i} \left(Y \mid \mathbf{X}_{\sim i} \right) \right) + \\ E_{\mathbf{X}_{\sim i}} \left(V_{X_i} \left(Y \mid \mathbf{X}_{\sim i} \right) \right) = V(Y)$$

If this = main effect on
non- X_i

$$E_{\mathbf{X}_{\sim i}} \left(V_{X_i} \left(Y | \mathbf{X}_{\sim i} \right) \right)$$





$$V_{\mathbf{X}_{\sim i}} \left(E_{X_i} \left(Y | \mathbf{X}_{\sim i} \right) \right)$$

... then all remaining variance must be due to X_i and its interactions; this divided the total variance gives the total effect T_i

How to compute S_i and T_i ?

If k is the number of factors we need $n(2 + k)$ simulation where n is the size of a ‘base’ matrix.


What is needed is a design (as in statistical DOE theory)

Building a Monte Carlo analysis

$$\begin{array}{ccc} x_{11} & x_{12} \dots & x_{1k} \\ x_{21} & x_{22} \dots & x_{2k} \\ \dots & \dots & \dots \\ x_{N1} & x_{N2} & x_{Nk1} \end{array}$$

Input matrix: each column is a sample of size N from the distribution of a factor

Each row is a sample trial of size k to generate a value of y



Examples of distributions of input factors

$$\mathbf{M}_1 = \begin{pmatrix} x_{11} & x_{12} & \dots & x_{1k} \\ x_{21} & x_{22} & \dots & x_{2k} \\ \dots & & & \\ x_{n1} & x_{n2} & \dots & x_{nk} \end{pmatrix}$$

Base matrix $\mathbf{M}_1$

$$\mathbf{M}_2 = \begin{pmatrix} x'_{11} & x'_{12} & \dots & x'_{1k} \\ x'_{21} & x'_{22} & \dots & x'_{2k} \\ \dots & & & \\ x'_{n1} & x'_{n2} & \dots & x'_{nk} \end{pmatrix}$$

Base matrix $\mathbf{M}_2$

$$\mathbf{N}_j = \begin{pmatrix} x'_{11} & x'_{12} & \dots & x_{1j} & \dots & x'_{1k} \\ x'_{21} & x'_{22} & \dots & x_{2j} & \dots & x'_{2k} \\ \dots & \dots & \dots & \dots & \dots & \dots \\ x'_{n1} & x'_{n2} & \dots & x_{nj} & \dots & x'_{nk} \end{pmatrix}$$

Matrix quasi- $\mathbf{M}_2$ for factor j

k of these are needed

'quasi' because all columns are from $\mathbf{M}_2$ but column j from $\mathbf{M}_1$

$$\mathbf{M}_1 = \begin{pmatrix} x_{11} & x_{12} & \dots & x_{1k} \\ x_{21} & x_{22} & \dots & x_{2k} \\ \dots & & & \\ x_{n1} & x_{n2} & \dots & x_{nk} \end{pmatrix}$$

Base matrix $\mathbf{M}_1$

$$\mathbf{M}_2 = \begin{pmatrix} x'_{11} & x'_{12} & \dots & x'_{1k} \\ x'_{21} & x'_{22} & \dots & x'_{2k} \\ \dots & & & \\ x'_{n1} & x'_{n2} & \dots & x'_{nk} \end{pmatrix}$$

Base matrix $\mathbf{M}_2$

S_j

T_j

$$\mathbf{N}_j = \begin{pmatrix} x'_{11} & x'_{12} & \dots & x_{1j} & \dots & x'_{1k} \\ x'_{21} & x'_{22} & \dots & x_{2j} & \dots & x'_{2k} \\ \dots & \dots & \dots & \dots & \dots & \dots \\ x'_{n1} & x'_{n2} & \dots & x_{nj} & \dots & x'_{nk} \end{pmatrix}$$

k of these are needed

Matrix quasi- $\mathbf{M}_2$ for factor j

How about Shapley?

$$S_i \leq \text{Sh}_i \leq T_i,$$

Shapley sits between S_i and T_i for uncorrelated input

Why?

Owen, A. B. (2014). "Sobol' indices and Shapley value."
SIAM/ASA Journal on Uncertainty Quantification, 2(1), 245–251.

$$S_i \leq Sh_i \leq T_i,$$

- S_i gets none of the interactions
- T_i gets all of the interactions belonging to X_i
- Shapley value splits each interaction term *equally* among the variables
- Shapley effects sum exactly to the total variance (or to 1 if normalised)

$T_i - S_i$ is the interaction that Shapley redistributes.



Shy



Leader



Diplomat

$$v(\text{Shy}) \text{ } 1 \text{ apple}$$

$$v(\text{Leader}) \text{ } 3 \text{ apples}$$

$$v(\text{Diplomat}) \text{ } 2 \text{ apples}$$

$$v(\text{Shy}, \text{Leader}) \text{ } 2 \text{ apples}$$

$$v(\text{Shy}, \text{Diplomat}) \text{ } 5 \text{ apples}$$

$$v(\text{Leader}, \text{Diplomat}) \text{ } 4 \text{ apples}$$

$$v(\text{Leader}, \text{Diplomat}, \text{Shy}) \text{ } 9 \text{ apples}$$

Shy and Leader have 2.33 apples each
 Diplomat has 3.333 apples


Images courtesy from
 Bertrand Iooss, Electricité de
 France (EDF)

A link between $I(X^j)$, a measure of permutation importance in ML, and the total sensitivity index

$$ST_j = \frac{I(X^j)}{2\text{Var}(Y)}$$

Antoniadis, Anestis, Sophie Lambert-Lacroix, and Jean-Michel Poggi. 2021. 'Random Forests for Global Sensitivity Analysis: A Selective Review'. Reliability Engineering & System Safety 206 (February): 107312. <https://doi.org/10.1016/j.ress.2020.107312>.

... but there are other methods that can be used for different settings, e.g. moment-independent:





Environmental Modelling & Software

Volume 34, June 2012, Pages 105-115




Model emulation and moment-independent sensitivity analysis: An application to environmental modelling

E. Borgonovo ^a, W. Castaings ^{b, c}, S. Tarantola ^d



Regionalised sensitivity analysis; the input distribution is split according to Y being **Behaviour** or **Non-Behaviour**

Exploring the multidimensional cube
(Go quasirandom)



An LP_τ sequence




Ilya M Sobol'
1926–2025

Sensitivity buffs use
quasirandom sequences,
preferentially Sobol' LP-TAU
sequences


Sobol' Sequences vrs Random numbers and regular grid

Slide courtesy of
Sergei Kucherenko,
Imperial College
London


Regular Grid/ 64 Points




Random Numbers/ 64 Points




Sobol Numbers/ 64 Points




Regular Grid/ 256 Points



Random Numbers/ 256 Points





Sobol Numbers/ 256 Points





Slide courtesy of Sergei
Kucherenko, Imperial
College >London



Projections of Different 2D sequences to 1D



Shadow effect
(64 -> 8 points)




Clustering





Uniform distribution

Part II – Who does GSA?



Rodrik's rules about models (and GSA)

- Economics is a collection of models; cherish their diversity.
- It's a model, not the model.
- Make your model simple enough to isolate specific causes and how they work, but not so simple that it leaves out **key interactions** among causes.
- Unrealistic assumptions are OK; unrealistic **critical assumptions** are not OK




→ When modelling, move horizontally rather than vertically

Exploratory modelling analysis (RAND corporation)

Explore scenarios and simulate



How wide is wide enough?



Jan H. Kwakkel and Erik Pruyt. Exploratory Modeling and Analysis, an approach for model-based foresight under deep uncertainty. Technological Forecasting and Social Change, 80(3):419–431, March 2013.

Application to Machine Learning

Example use SA to ascertain that an algorithm does not make implicit use of protected attributes

More reading about explainable ML

Lu, Xuefei, and Emanuele Borgonovo. 2026. "How Do I Explain That Decision? A New Approach Combining Counterfactuals and Sensitivity Analyses." *European Journal of Operational Research* 334 (1): 200–215. <https://doi.org/10.1016/j.ejor.2026.04.035>.

Giulia Vannucci, Roberta Siciliano, Andrea Saltelli, 2026, Global sensitivity analysis in random forests: unveiling generative variable importance. *Statistical Methods and Applications*, 34(5), <https://doi.org/10.1007/s10260-026-00839-y>.

Bénese, C., Gamboa, F., Loubes, J.-M., & Boissin, T. (2022). Fairness seen as Global Sensitivity Analysis. *Machine Learning*. <https://doi.org/10.1007/s10994-022-06202-y>

PROTECTED ATTRIBUTES:

- Age
- Disability
- National Origin
- Race/color
- Religion
- Sex
- (From the US Equal Opportunity Employment Commission)

More reading about explainable ML

Bénesse, C., Gamboa, F., Loubes, J.-M., & Boissin, T. (2022). Fairness seen as Global Sensitivity Analysis. *Machine Learning*. <https://doi.org/10.1007/s10994-022-06202-y>

“fairness can be defined using a special framework based on Global Sensitivity Analysis ...”

[Find a journal](#) [Publish with us](#) [Track your research](#) |

[Saved research](#) [Cart](#)

[Home](#) > [Machine Learning](#) > [Article](#)


Fairness seen as global sensitivity analysis

Published: 24 June 2022

Volume 113, pages 3205–3232 (2024) [Cite this article](#)

[Download PDF](#) 

 [Save article](#)



Machine Learning

[Aims and scope](#) →

[Submit manuscript](#) →

[Clément Bénesse](#) , [Fabrice Gamboa](#), [Jean-Michel Loubes](#) & [Thibaut Boissin](#)

Part of a collection:
[Special Issue on Safe and Fair Machine Learning](#)

More reading about explainable ML: combining counterfactual and Shapley values

Lu, Xuefei, and Emanuele Borgonovo. 2026. "How Do I Explain That Decision? A New Approach Combining Counterfactuals and Sensitivity Analyses." *European Journal of Operational Research* 334 (1): 200–215.
<https://doi.org/10.1016/j.ejor.2026.04.035>.

"In machine learning, [the idea is]to treat features as players and model predictions as rewards"



The screenshot shows the ScienceDirect article page for the paper "How do I explain that decision? A new approach combining counterfactuals and sensitivity analyses" by Xuefei Lu and Emanuele Borgonovo. The page includes the ScienceDirect logo, the journal title "European Journal of Operational Research", and the article title. It also displays the authors' names, a "Show more" link, and various sharing options like "Cite", "Add to Mendeley", and "Share". The DOI "10.1016/j.ejor.2026.04.035" is visible. At the bottom, there is a blue button for "Access through Universitat Pompeu Fabra" and links for "Purchase PDF" and "Access through another organization".

ScienceDirect Journals & Books Help Search Andrea Saltelli

Research article Get rights and content »

Decision Support

How do I explain that decision? A new approach combining counterfactuals and sensitivity analyses


Xuefei Lu ^a, Emanuele Borgonovo ^{b,c} Show more ▾

Cite Add to Mendeley Share 10.1016/j.ejor.2026.04.035 »

European Journal of Operational Research

Access through Universitat Pompeu Fabra Purchase PDF Access through another organization

Sensitivity analysis in the garden



“Will different researchers [73 teams] converge on similar findings when analysing the same data?”

“...teams’ results varied greatly, ranging from large negative to large positive effects”

PNAS

RESEARCH ARTICLE | SOCIAL SCIENCES

OPEN




Observing many researchers using the same data and hypothesis reveals a hidden universe of uncertainty

Edited by Douglas Massey, Princeton University, Princeton, NJ; received March 6, 2022; accepted August 22, 2022

Simulate different paths using triggers

k_i	Variable	Description
1	<i>age</i>	Inclusion or non-inclusion of the covariates age and square age of participants
2	<i>all – ethnic</i>	Inclusion or non-inclusion of a covariate that takes into account all ethnic categories
3	<i>bootstrap</i>	Inclusion or non-inclusion of the bootstrap procedure
4	<i>cont – europe</i>	Inclusion or non-inclusion of continental Europe countries in the analysis
5	<i>income</i>	Inclusion or non-inclusion of the covariate related to the income
3	<i>bootstrap</i>	Inclusion or non-inclusion of the bootstrap procedure
4	<i>cont – europe</i>	Inclusion or non-inclusion of continental Europe countries in the analysis
9	<i>income</i>	Inclusion or non-inclusion of the covariate related to the income
10	<i>pop – wb</i>	Inclusion or non-inclusion of the covariate related to the information about country-specific numerosity of population
11	<i>rich – dems</i>	Inclusion or non-inclusion of rich democracies in the analysis
12	<i>SE</i>	Inclusion or non-inclusion of clustered standard error
13	<i>socx – oecd</i>	Inclusion or non-inclusion of the covariate related to social expenditure
14	<i>X1</i>	Inclusion or non-inclusion of the covariates related to the migration indexes
15	<i>Y</i>	Choice of policy dependent variable

Table 1. Descriptions of variables included in the analysis.



Our uncertainty analysis broadly reproduces the results of the multi-analyst experiment


Our sensitivity analysis explains where the missing variance went: **interactions**

Part III GSA's dos and don'ts


➔ Don't sample just parameters and boundary conditions: explore the space of assumptions thoroughly

One can sample more than just factors:

- modelling assumptions,
- alternative data sets,
- resolution levels,
- policies



Look at policy options versus existing uncertainties: e.g., test whether the choice of the model determines the choice of the policy



Open Access


November 2000

Sensitivity Analysis as an Ingredient of Modeling

F. Campolongo, A. Saltelli, S. Tarantola


Statist. Sci. 15(4): 377-395 (November 2000). DOI: 10.1214/ss/1009213004

Source: Saltelli, A. (2019). Statistical versus mathematical modelling: A short comment. Nature Communications, 10, 1–3.
<https://doi.org/10.1038/s41467-019-11865-8>



Don't confuse the map with the territory


If you do, sensitivity analysis will not save you




<<...If the model is wrong or if it is a poor representation of reality, determining the sensitivity of an individual parameter in the model is a meaningless pursuit>>

One of the examples discussed concerns the **Yucca Mountain** repository for radioactive waste. TSPA model (for total system performance assessment) for safety analysis.


TSPA is Composed of 286 sub-models.





TSPA (like any other model) **relies on assumptions** → one is the low permeability of the geological formation → long time for the water to percolate from surface to disposal





“Scientific mathematical modelling should involve constant efforts to falsify the model”

➔ Organized skepticism (as per CUDOS)

Communalism, Universalism, Disinterestedness, Organized Skepticism, from sociology of science, Robert K. Merton.

Don't run the model just once

There is much to learn by running the model a few times, especially during model building

Lubarsky's Law of Cybernetic Entomology:
there is always one more bug!



A model routinely used to produce point estimates may become non-conservative when the uncertainty is plugged in

In a numerical experiment relating to a real-life application, the range of uncertainty of each input is a crucial input to the analysis, and often the most expensive to get